

## Case Report

# Subcutaneous Granulomatous Inflammation due to Basidiobolomycosis: Case Reports of 3 Patients in Buruli Ulcer Endemic Areas in Benin

Luc V. C. Brun <sup>1</sup>, Jean Jacques Roux,<sup>2</sup> Ghislain E. Sopoh <sup>3,4</sup>,  
Julia Aguiar,<sup>5</sup> Miriam Eddyani,<sup>6</sup> Wayne M. Meyers,<sup>7</sup> Dirk Stubbe,<sup>8</sup> Marie T. Akele Akpo,<sup>9</sup>  
Françoise Portaels,<sup>6</sup> and Bouke C. de Jong<sup>6</sup>

<sup>1</sup>Department of Pathology, School of Medicine, University of Parakou, 03 BP 333 Parakou, Benin

<sup>2</sup>Department of Pathology, Hospital of Chambéry, Place Lucien Biset, 73000 Chambéry, France

<sup>3</sup>Buruli Ulcer Treatment Center, Allada, Benin

<sup>4</sup>Regional Institute of Public Health, Ouidah, Benin

<sup>5</sup>Nutritional Center of Gbemontin, Zagnanado, Benin

<sup>6</sup>Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerpen, Belgium

<sup>7</sup>Armed Forces Institute of Pathology, Washington, DC, USA

<sup>8</sup>BCCM/IHEM Biomedical Fungi and Yeasts Collection, Scientific Institute of Public Health, Brussels, Belgium

<sup>9</sup>Department of Pathology, School of Medicine, University of Abomey Calavi, Cotonou, Benin

Correspondence should be addressed to Luc V. C. Brun; [luc.brun2013@gmail.com](mailto:luc.brun2013@gmail.com)

Received 11 August 2017; Accepted 11 December 2017; Published 10 January 2018

Academic Editor: Mark Li-cheng Wu

Copyright © 2018 Luc V. C. Brun et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Background.** Basidiobolomycosis is a rare subcutaneous mycosis, which can be mistaken for several other diseases, such as soft tissue tumors, lymphoma, or Buruli ulcer in the preulcerative stage. Microbiological confirmation by PCR for *Basidiobolus ranarum* and culture yield the most specific diagnosis, yet they are not widely available in endemic areas and with varying sensitivity. A combination of histopathological findings, namely, granulomatous inflammation with giant cells, septate hyphal fragments, and the Splendore-Hoeppli phenomenon, can confirm basidiobolomycosis in patients presenting with painless, hard induration of soft tissue. **Case Presentations.** We report on three patients misdiagnosed as suffering from Buruli ulcer, who did not respond to Buruli treatment. Histopathological review of the tissue sections from these patients suggests basidiobolomycosis. All patients had been lost to follow-up, and none received antifungal therapy. On visiting the patients at their homes, two were reported to have died of unknown causes. The third patient was found alive and well and had experienced local spontaneous healing. **Conclusion.** Basidiobolomycosis is a rare subcutaneous fungal disease mimicking preulcerative Buruli ulcer. We stress the importance of the early recognition by clinicians and pathologists of this treatable disease, so patients can timely receive antifungal therapy.

## 1. Introduction

Basidiobolomycosis is a subcutaneous mycosis that occurs in tropical and subtropical regions of Asia, Africa, and South America, caused by *Basidiobolus ranarum* Eidam or related species [1]. The disease mainly affects children and tends to preferentially affect the thighs, buttocks, or trunks. The clinical and histopathological signs at first do not evoke a fungal disease. The initial infection is subacute, sometimes

marked by a pseudophlegmon or erysipelas. The evolution is slow and gradual, with periods of remissions over several months or years, in the form of an extensive plaque. Patients on immunosuppressive therapy may be at higher risk of developing progressive disease [2]. Clinical examination reveals inhomogeneous swellings. Palpation reveals irregular firm nodules. The edema is nonpitting and hard as wood, and the skin is rigid and cannot wrinkle. The symptomatology varies, depending on the evolutionary stage (early versus

chronic infection). In early infection, nodules are hot and slightly painful. In chronic infection, they are cold and painless. This soft tissue infection usually respects the joints and muscles, infrequently causing functional impairment [3].

Although medical treatment by oral potassium iodide and ketoconazole can be effective [4], the untreated infection can be fatal. Over the past 40 years, 179 cases of subcutaneous basidiobolomycosis have been reported worldwide [5], including West Africa [6–10]. Among them, there were only seven cases, recently reported in Benin [10].

*Basidiobolus ranarum* belongs to the fungal phylum Entomophthoromycota. This phylum holds over 250 species, mostly soil or litter saprophytes, but also numerous arthropod pathogens. It has emerged as one of the major lineages of the dissolved phylum Zygomycota (which, at the time, also included the now independent subphylum Mucoromycotina). Within the phylum Entomophthoromycota, two species are known as human pathogens: *Conidiobolus coronatus* (Costantin) Batko (in the order Entomophthorales, causing rhinophycomycosis) and *B. ranarum* (in the order Basidiobolales). The genus *Basidiobolus* contains at least six species [11]. In addition to *B. ranarum*, also *B. meristosporus* Drechsler and *B. haptosporus* Drechsler have been identified as causal agents of basidiobolomycosis. These latter two names are currently accepted as synonyms of *B. ranarum*. However, this is not corroborated by the multigene analysis of Gryganskyi [11] and more taxonomic research will be needed to resolve this delineation of the taxonomy. *Basidiobolus* spp. have been isolated from reptile or amphibian dung, but also bats and other (insectivorous) animals have been identified as carriers and possible reservoirs [12–15].

The transmission route of *Basidiobolus* spp. is poorly understood. Basidiobolomycosis develops in the subcutaneous soft tissue and yet can involve muscles and surrounding organs, as well as gastrointestinal lesions [16, 17].

Basidiobolomycosis can be mistaken for Buruli ulcer (BU), an infectious skin disease due to *Mycobacterium ulcerans*, also endemic in some countries in Africa, Asia, South America, Mexico, and Australia, which presents as a painless papule, nodule, plaque, edema, or ulcer. The infrequent encounter of basidiobolomycosis and its symptoms that can mimic a BU plaque or oedematous form limits clinician experience in its recognition in the differential diagnosis. Histopathological examination of a tissue biopsy can provide diagnostic clues, when a dermohypodermic granuloma is observed, rich in eosinophils and giant cells, with the presence of hyphae. The phenomenon of Splendore-Hoeppli consists of an asteroid body, corresponding to eosinophilic material often in a stellate or club form [18], organized around various microorganisms such as bacteria, fungi, parasites, or even biologically inert substances [18], yet mostly associated with basidiobolomycosis [4, 19]. Microbiological confirmation of the diagnosis of basidiobolomycosis is based on the detection of colonies of *B. ranarum* after seeding a biopsy specimen on Sabouraud medium [20] or PCR detection of specific targets. We describe three patients with subcutaneous basidiobolomycosis in a Buruli Treatment Center (CDTUB) in Benin, two of whom had died on follow-up.

## 2. Case Presentation

**2.1. Patient 1.** A malnourished three-year-old girl living in a BU-endemic area was admitted to the CDTUB of Zagnanado in 1997 for a two-month history of swelling of the right flank. The lesion was a hard painless plaque. Based on the clinical signs, in conjunction with residence in a BU-endemic area, an initial diagnosis of BU was made. Surgical excision was performed and samples were collected for bacteriology, *M. ulcerans* specific PCR, and histopathology. Treatment with cloxacillin and gentamycin was instituted. Direct microscopy of the tissue samples after Ziehl-Neelsen staining was negative for acid fast bacilli (AFB). The IS2404 PCR for *M. ulcerans* was also negative. However the histopathological examination of the tissue samples after Hematoxylin-Eosin (HE) and Gomori-Methenamine-Silver (GMS) staining provided strong support for the diagnosis of basidiobolomycosis. PCR for the detection of *B. ranarum* DNA was not performed. No information was available about the disease progression in this patient at the time she was lost to follow-up. However, she was recently reidentified and had undergone spontaneous healing.

**2.2. Patient 2.** A 43-year-old man living in a rural BU-endemic area presented in the CDTUB of Zagnanado in 2009 with an extensive atypical cutaneous plaque on the right thigh that had developed over six months. He received traditional herbal treatment before he presented. On palpation, the lesion was painless with very hard induration, and the patient was afebrile. Based on the clinical signs and symptoms, as well as residence in a BU-endemic region, an initial diagnosis of BU was made. Surgical excision was performed and tissue samples were collected for laboratory confirmation of BU. Laboratory tests (direct smear examination after Ziehl-Neelsen staining, culture, and IS2404 PCR) were negative.

Histopathological examinations revealed a granulomatous inflammation with giant cells without caseating necrosis, reported as tuberculoïd granulomas. The patient had received antibiotic treatment for BU (streptomycin and rifampicin for 8 weeks) and cloxacillin, without improvement.

Subsequently, the lesion extended towards the right leg and buttock. The patient was lost to follow-up. His relatives mentioned that the lesions progressed from nonulcerated lesions to an ulcerated one. He died of unknown causes.

Five years after the patient's death, histopathological reexamination suggested the diagnosis of basidiobolomycosis based on the presence of eosinophilic and giant multinucleated cells mixed with septate hyphal fragments and the Splendore-Hoeppli phenomenon. We submitted frozen tissue samples for PCR at the BCCM/IHEM culture collection for biomedical fungi at the Scientific Institute of Public Health in Belgium, but these were negative for *B. ranarum*.

**2.3. Patient 3.** A three-year-old boy living in a rural BU-endemic area was admitted in the CDTUB of Zagnanado in 2012, for an atypical cutaneous and extensive plaque-like lesion. The lesion was located on the right thigh (Figure 1(a)) and clinically resembled BU or a soft tissue tumor. The lesion was painless, had the hardness of wood, and started one



FIGURE 1: Clinical aspects (patient 3). (a) Lesion on R thigh, recurrence after initial resection. (b) The same patient three years later (a few months before his death).

month before admission. The patient was in good overall physical condition. Laboratory tests on biopsy samples for BU (direct smear examination after Ziehl-Neelsen staining, culture, and IS2404 PCR) were negative.

The initial histopathological examination was interpreted as “possible BU” because of the observed granulomatous inflammation. Despite BU specific antibiotic treatment (streptomycin and rifampicin for 8 weeks) and cloxacillin, as well as surgical excision of the lesion, the induration recurred before the surgical wound had healed. The lesion then extended towards the leg and buttock. The patient was lost to follow-up, despite attempts to visit him at home, and had been taken to a traditional practitioner. When the histopathological diagnosis was revised as suggesting basidiobolomycosis rather than BU, the parents refused free treatment with ketoconazole. The lesion extended to the whole right lower limb with a pseudoelephantiasis aspect (Figure 1(b)), and the skin ulcerated on the calf. The patient died two years after the onset of the illness. A PCR for *B. ranarum*, performed at BCCM/IHEM on DNA extracted from frozen tissue suspensions, was negative.

Histopathological features were consistent with basidiobolomycosis, based on the presence of eosinophilic cells and giant multinucleated cells mixed with rare septate hyphal fragments and Splendore-Hoeppli material.

The families of all three patients provided written informed consent for these anonymized case reports despite refusal of free treatment (patient 3).

**2.4. Pathology Results.** Histopathological results after HE staining of all three cases showed tuberculoïd granulomata with giant cells (Figure 2(a)), numerous lymphocytes, histiocytes and eosinophilic cells (Figure 2(b)), an amorphous eosinophilic material also known as the “Splendore-Hoeppli phenomenon” (Figure 2(c)), and 10  $\mu\text{m}$  diameter septate hyphal fragments (Figure 2(b)), confirmed also by the Gomori-Grocott staining (Figure 2(d)). These hyphal fragments sometimes appeared centered within a Splendore-Hoeppli phenomenon material or were found in the cytoplasm of multinucleated giant cells.

### 3. Discussion

In these three patients the diagnosis of basidiobolomycosis was histologically suggested, based on the presence of a very hard panniculitis with a significant sclero-inflammatory response (lymphocytes, histiocytes, eosinophiles, and giant cells), few septate hyphae, and the presence of the Splendore-Hoeppli phenomenon. PCR was negative for *B. ranarum*. The differential diagnosis provided for basidiobolomycosis includes soft tissue tumors, such as a synovial sarcoma [21], Hodgkin lymphoma [22], and mycetoma and BU [5]. Indeed, a tuberculoïd granuloma with giant cells can also be observed in BU lesions, especially on healing. However, neither the septate hyphal fragments nor the Splendore-Hoeppli phenomenon is observed in BU lesions.

Importantly, BU differs from basidiobolomycosis in etiology and management. The Splendore-Hoeppli phenomenon itself is not specific for basidiobolomycosis [18] and can also be observed in other infections, such as botryomycosis [23] bronchocentric granulomatosis due to *Aspergillus* [24] mycetoma [25] and *Pityrosporum* folliculitis skin infection [26], although the combination of the clinical presentation, the Splendore-Hoeppli phenomenon, and septate hyphae suggests basidiobolomycosis. In addition to a predilection for lungs rather than the subcutis, *Aspergillus* can be recognized by septate hyphae that branch at 45° angles and by vascular invasion, while mycetoma produces grains and has a very distinct clinical presentation from basidiobolomycosis. Since culture of the offending fungus *B. ranarum* is difficult [3], clinical and histopathological features can help to suggest the diagnosis of basidiobolomycosis. Among seven cases recently reported in Benin by dermatologists, five whose tissues were sampled by biopsy were diagnosed using histopathology [10]. Early recognition of basidiobolomycosis would allow early effective treatment, even in the absence of culture or PCR positive results. The detection of fungal pathogens by PCR is particularly challenging. Fungal cell walls are not easily lysed for the release of DNA, which leads to false-negative PCR results [27]. Isolation of *Basidiobolus ranarum* DNA from archival formalin-fixed, paraffin embedded (FFPE) tissue




FIGURE 2: Histopathology of basidiobolomycosis. (a) Granulomatous inflammation with eosinophils and multinucleated giant cells in the subcutaneous tissue. Hematoxylin-Eosin, 50x. (b) Granulomatous inflammation with eosinophils cells and septate hyphal fragments. Hematoxylin-Eosin, 100x. (c) Granulomatous inflammation with eosinophils, multinucleated giant cells, and the Splendore-Hoeppli phenomenon. Hematoxylin-Eosin, 100x. (d) Septate hyphal fragments (arrow). Gomori-Grocott staining, 400x.

blocks has been reported, with a protocol allowing reliable purification of fungal DNA [28, 29]. The antifungals that have most been used for treatment of basidiobolomycosis are oral itraconazole [10, 30, 31] and oral ketoconazole at a dose of 400 mg per os once daily [32] or 7 to 10 mg per kilogram once daily (for children), for two to six months [6, 7, 10, 33]; posaconazole has also been effectively used in one case report [34]. Potassium iodide may be another effective treatment option [35].

None of the patients in our series benefited from ketoconazole treatment. The third patient's parents refused this outpatient treatment, and his disease progressed, causing elephantiasis-like lymphedema (Figure 1(b)), as described by Kamalam and Thambiah [36]. We hypothesize that the parents' refusal stemmed from a loss of confidence in allopathic medicine due to the delayed diagnosis of basidiobolomycosis, which was preceded by erroneous treatment for BU including surgical excision. With the failure of allopathic medical and surgical therapies the parents interpreted the unfavorable evolution as evidence for witchcraft (enchantment); in Benin this culturally supported framework for understanding disease warrants traditional therapies including herbal treatments, rather than allopathic approaches [37–40]. It is important that health workers, especially in BU-endemic areas, recognize the clinical symptoms of basidiobolomycosis and ask for histological testing in order to avoid delays

in diagnosis and appropriate treatment for this potentially devastating disease.

#### 4. Conclusion

Clinicians need to recognize wood-like induration of skin and soft tissue as a potential deep-seated fungal infection, also in BU-endemic areas. Such patients first and foremost need antifungal treatment, preferably after a biopsy is taken for histopathological features characteristic for basidiobolomycosis, rather than a wide excision.

Failure to recognize this clinical entity may result in therapeutic failure and may have contributed to the death of two of these patients. Clinical instruction on this presentation is therefore of paramount importance in tropical healthcare education.

#### Consent

Written informed consent was obtained from the families of the patients for publication of this case report and any accompanying images.

#### Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Authors' Contributions

Luc V. C. Brun and Bouke C. de Jong conceived the manuscript. Luc V. C. Brun compiled the patient information. Luc V. C. Brun and Bouke C. de Jong wrote the manuscript. Luc V. C. Brun, Wayne M. Meyers, Marie T. Akele Akpo, and Jean Jacques Roux read the histopathological slides. Françoise Portaels and Miriam Eddyani participated in the design of the study and the PCR for Buruli ulcer on skin biopsies. Dirk Stubbe participated in the PCR for basidiobolomycosis diagnosis. Ghislain E. Sopoh and Julia Aguiar participated in making the skin biopsy of the patients. Bouke C. de Jong participated in its design and coordination, assisted in the PCR for Buruli ulcer, and helped in drafting the manuscript. All authors read and approved the final manuscript.





## Acknowledgments

This research was supported by the UBS Optimus and the Medicor Foundations, as well as the SOFI funds from the EWI Ministry and ERC starting Grant 311725 INTERRUPTB to Bouke C. de Jong. The authors are grateful to the families of the patients, who agreed on the publication of this case report. They are also grateful to Paulin Déguénon and Michel Assogbaga from the Nutritional health center "Gbemontin" of Zagnanado and to Thierry Kinkpé for their support.

## References

- [1] J. Coremans-Pelseeneer, "Epidemiology of basidiobolomycosis," *Ann Soc Belge Med Trop*, vol. 52, pp. 315–328, 1972.
- [2] P. Sethi, D. Balakrishnan, S. Surendran, and Z. U. Mohamed, "Fulminant zygomycosis of graft liver following liver transplantation," *BMJ Case Reports*, vol. 2016, Article ID 214097, 2016.
- [3] A. Basset, "Clinical and treatment aspects of Basidiobolomycosis," *Ann Soc Belge Med Trop*, vol. 52, pp. 307–314, 1972.
- [4] L. Mendoza, R. Vilela, K. Voelz, A. S. Ibrahim, K. Voigt, and S. C. Lee, "Human fungal pathogens of Mucorales and Entomophthorales," *Cold Spring Harbor Perspectives in Medicine*, vol. 5, no. 4, Article ID a019562, 2015.
- [5] K. Kombaté, B. Saka, and A. Mouhari-Toure, "Basidiobolomycosis: a review," *Med Santé Trop*, vol. 22, pp. 145–152, 2012.
- [6] B. Saka, K. Kombaté, A. Mouhari-Toure et al., "Probable basidiobolomycosis in a Togolese rural young successfully treated with ketoconazole," *Bulletin de la Société de Pathologie Exotique*, vol. 103, no. 5, pp. 293–295, 2010.
- [7] A. Sangare, P. Yoboue, and V. Bamba, "Basidiobolomycosis: some aspects, about 10 cases observed in eleven years in the dermato-venerology clinic of the university hospital of Treichville," *Med d'Afr Noire*, vol. 47, pp. 8–9, 2000.
- [8] F. Atadokpede, H. Adégbidi, and J. Gnossiké, "Epidemiological and clinical aspects of the entomophthoromycosis in Pobé, Bénin," *Ann Dermatol Vénérol*, vol. 142, 2015.
- [9] "Two beninese entomophthoromycosis (basidiobolomycosis et conidiobolomycosis) cases diagnosed in the university hospital of Angers in 2010," *Journal de Mycologie Médicale*, vol. 21, no. 3, pp. 221–235, 2011, Résumés des conférences et des communications du congrès de la Société française de mycologie médicale, Strasbourg, France, 18–19 mai 2011.
- [10] F. Atadokpédé, J. Gnossiké, H. Adégbidi et al., "Cutaneous basidiobolomycosis: Seven cases in southern Benin," *Annales de Dermatologie et de Venereologie*, vol. 144, no. 4, pp. 250–254, 2017.
- [11] A. P. Gryganskyi, R. A. Humber, M. E. Smith et al., "Phylogenetic lineages in Entomophthoromycota," *Persoonia*, vol. 30, pp. 94–105, 2013.
- [12] R. Speare and A. D. Thomas, "Kangaroos and wallabies as carriers of Basidiobolus haptosporus," *Australian Veterinary Journal*, vol. 62, no. 6, pp. 209–210, 1985.
- [13] H. C. Gugnani and J. I. Okafor, "Mycotic flora of the intestine and other internal organs of certain reptiles and amphibians with special reference to characterization of Basidiobolus isolates," *Mykosen*, vol. 23, no. 5, pp. 260–268, 1980.
- [14] V. P. Chaturvedi, H. S. Randhawa, Z. U. Khan, N. Singh, and S. Kini, "Prevalence of Basidiobolus ranarum Eidam in the intestinal tract of an insectivorous bat, *Rhinopoma hardwickei* hardwickei Gray, in Delhi," *Medical Mycology*, vol. 22, no. 3, pp. 185–189, 1984.
- [15] R. T. Nelson, D. TeStrake, and B. J. Cochrane, "The distribution of Basidiobolus and Conidiobolus in soil and litter at sites near Tampa, Florida," *Mycologia*, vol. 90, no. 5, pp. 761–766, 1998.
- [16] T. M. Pasha, J. A. Leighton, J. D. Smilack, J. Heppell, T. V. Colby, and L. Kaufman, "Basidiobolomycosis: An unusual fungal infection mimicking inflammatory bowel disease," *Gastroenterology*, vol. 112, no. 1, pp. 250–254, 1997.
- [17] M. El-Shabrawi, L. Madkour, and N. Voigt, "Pediatric Invasive Gastrointestinal Fungal Infections: Causative Agents and Diagnostic Modalities," *Microbiology Research Journal International*, vol. 19, no. 2, pp. 1–11, 2017.
- [18] M. R. Hussein, "Mucocutaneous Splendore-Hoepli phenomenon," *Journal of Cutaneous Pathology*, vol. 35, no. 11, pp. 979–988, 2008.
- [19] J. E. Rothhardt, V. U. Schwartz, and K. Voigt, "Entomophthorales," in *Molecular Detection of Human Fungal Pathogens*, L. Dongyou, Ed., chapter 83, pp. 723–734, Taylor & Francis CRC Press, 1st edition, 2011.
- [20] Z. U. Khan, M. Khourshed, R. Makar et al., "Basidiobolus ranarum as an etiologic agent of gastrointestinal zygomycosis," *Journal of Clinical Microbiology*, vol. 39, no. 6, pp. 2360–2363, 2001.
- [21] Sivaraman, D. M. Thappa, Karthikeyan, and Hemanthkumar, "Subcutaneous phycomycosis mimicking synovial sarcoma," *International Journal of Dermatology*, vol. 38, no. 12, pp. 920–923, 1999.
- [22] A. L. Bittencourt, G. Serra, M. Sadigursky, M. G. Araujo, M. C. Campos, and L. C. Sampaio, "Subcutaneous zygomycosis caused by Basidiobolus haptosporus: Presentation of a case mimicking Burkitt's lymphoma," *The American Journal of Tropical Medicine and Hygiene*, vol. 31, no. 2, pp. 370–373, 1982.
- [23] H. Marques Barreiros, H. Cunha, and E. Bártolo, "Botryomycosis in an immunocompetent woman," *Journal of Dermatological Case Reports*, vol. 7, no. 1, pp. 29–30, 2013.
- [24] Y. Yoshikawa, L. D. Truong, and T. Watanabe, "Splendore-Hoepli phenomenon in bronchocentric granulomatosis," *Thorax*, vol. 43, no. 2, pp. 157–158, 1988.
- [25] J. W. Kang, J. S. Ku, C.-H. Kim, and J.-H. Yoon, "Intranasal mycetoma-induced Splendore-Hoepli phenomenon," *Otolaryngology—Head and Neck Surgery*, vol. 142, no. 3, pp. 456–457, 2010.
- [26] O. J. Clemmensen and H. Hagdrup, "Splendore-Hoepli phenomenon in Pityrosporum folliculitis (pseudactinomycosis of the skin)," *Journal of Cutaneous Pathology*, vol. 18, no. 4, pp. 293–297, 1991.

- [27] P. D. Khot and D. N. Fredrick, "PCR-based diagnosis of human fungal infections," *Expert Review of Anti-infective Therapy*, vol. 7, no. 10, pp. 1201–1221, 2009.
- [28] H. F. M. El-Shabrawi, N. M. Kamal, R. Jouini, A. Al-Harbi, K. Voigt, and T. Al-Malki, "Gastrointestinal basidiobolomycosis: An emerging fungal infection causing bowel perforation in a child," *Journal of Medical Microbiology*, vol. 60, no. 9, pp. 1395–1402, 2011.
- [29] M. H. El-Shabrawi, N. M. Kamal, K. Kaerger, and K. Voigt, "Diagnosis of gastrointestinal basidiobolomycosis: A mini-review," *Mycoses*, vol. 57, no. 3, pp. 138–143, 2014.
- [30] A. Mahé, M. Huerre, S. Kéita, F. Traoré, and P. Bobin, "Subcutaneous phycomycosis successfully treated with itraconazole," *Annales de Dermatologie et de Venerologie*, vol. 123, no. 3, pp. 182–184, 1996.
- [31] R. Mathew, S. Kumaravel, S. Kuruvilla et al., "Successful treatment of extensive basidiobolomycosis with oral itraconazole in a child," *International Journal of Dermatology*, vol. 44, no. 7, pp. 572–575, 2005.
- [32] L. M. Vianna, M. V. Lacerda, and M. A. Moraes, "Case report of subcutaneous entomophthoromycosis with retroperitoneal invasion," *Journal of the Brazilian Society of Tropical Medicine*, vol. 38, no. 4, pp. 348–350, 2005.
- [33] J. M. Lopes Costa, S. G. Marques, L. N. Barbosa et al., "Use of ketoconazole for treating mucocutaneous entomophthoromycosis: A case report," *Anais Brasileiros de Dermatologia*, vol. 79, no. 3, pp. 329–334, 2004.
- [34] S. R. Rose, M. D. Lindsley, S. F. Hurst, C. D. Paddock, T. Damodaran, and J. Bennett, "Gastrointestinal basidiobolomycosis treated with posaconazole," *Medical Mycology Case Reports*, vol. 2, pp. 11–14, 2013.
- [35] A. Sanaei Dashti, A. Nasimfar, H. Hosseini Khorami et al., "Gastro-intestinal basidiobolomycosis in a 2-year-old boy: dramatic response to potassium iodide," *Paediatrics and International Child Health*, pp. 1–4, 2016.
- [36] A. Kamalam and A. S. Thambiah, "Lymphedema and elephantiasis in Basidiobolomycosis," *Mycosen*, vol. 25, pp. 508–511, 1982.
- [37] I. Aujoulat, C. Johnson, C. Zinsou, A. Guédénon, and F. Portaels, "Psychosocial aspects of health seeking behaviours of patients with Buruli ulcer in southern Benin," *Tropical Medicine & International Health*, vol. 8, no. 8, pp. 750–759, 2003.
- [38] R. C. Johnson, M. Makoutode, and R. Hougnihin, "Le traitement traditionnel de l'ulcère de Buruli au Bénin," *Med Trop*, vol. 64, pp. 145–150, 2004.
- [39] K. Kibadi, M. Boelaert, M. Kayinua et al., "Therapeutic itineraries of patients with ulcerated forms of *Mycobacterium ulcerans* (Buruli ulcer) disease in a rural health zone in the democratic Republic of Congo," *Tropical Medicine & International Health*, vol. 14, no. 9, pp. 1110–1116, 2009.
- [40] P. Charlier, L. Brun, G. L. de la Grandmaison, and C. Hervé, "Medical students from Parakou (Benin) and West-African traditional beliefs on death and cadavers," *African Health Sciences*, vol. 12, no. 4, pp. 443–445, 2012.



**Hindawi**  
Submit your manuscripts at  
<https://www.hindawi.com>

